bs-20812R

[Primary Antibody]

BIOSS ANTIBODIES

Www.bioss.com.cn

sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

ZP2 Rabbit pAb

Target: ZP2

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: The zona pellucida is an extracellular matrix that surrounds the oocyte and early embryo. It is composed of three glycoproteins

with various functions during fertilization and preimplantation development. The glycosylated mature peptide is one of the structural components of the zona pellucida and functions in secondary binding and penetration of acrosome-reacted spermatozoa. Female mice lacking this gene do not form a stable zona matrix and are sterile. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Feb 2014]

Applications: IHC-P (1:100-500)


IHC-F (1:100-500) IF (1:100-500)

Reactivity: Mouse, Rat


Predicted MW.: ^{76 kDa}

Subcellular Location: Secreted

- VALIDATION IMAGES -



Paraformaldehyde-fixed, paraffin embedded (Mouse placenta); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Processed zona pellucida sperm-binding protein 2; ZP2) Polyclonal Antibody, Unconjugated (bs-20812R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat ovary); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Processed zona pellucida sperm-binding protein 2; ZP2) Polyclonal Antibody, Unconjugated (bs-20812R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

- SELECTED CITATIONS -

• [IF=3.777] Ling Shuai. et al. Hydroxychloroquine protects against autoimmune premature ovarian insufficiency by modulating the Treg/Th17 cell ratio in BALB/c mice. AM J REPROD IMMUNOL. 2023 Feb;: IHC; Mouse. 36752682