

bsm-60616R

[Primary Antibody]

SF1 Recombinant Rabbit mAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit

Clonality: Recombinant

Target: SF1

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene encodes a nuclear pre-mRNA splicing factor. The encoded protein specifically recognizes the intron branch point sequence and is required for the early stages of spliceosome assembly. Alternate splicing results in multiple transcript variants. [provided by RefSeq, May 2010]

Isotype: IgG

CloneNo.: E1F12

Applications: WB (1:500-2000)

IHC-P (1:100-500)

IHC-F (1:100-500)

IF (1:100-500)


Flow-Cyt (1:50-100)

ICC/IF (1:50-200)


Reactivity: Human, Mouse, Rat

Subcellular Location: Nucleus


— VALIDATION IMAGES —



25 ug total protein per lane of various lysates (see on figure) probed with SF1 monoclonal antibody, unconjugated (bsm-60616R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Rat Stomach; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SF1 Monoclonal Antibody, Unconjugated (bsm-60616R) at 1:150 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Mouse Stomach; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with SF1 Monoclonal Antibody, Unconjugated (bsm-60616R) at 1:150 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.