

bs-2756R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


CYLD Rabbit pAb**— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 1540**SWISS:** Q9NQC7**Target:** CYLD**Immunogen:** KLH conjugated synthetic peptide derived from human cylindromatosis 1: 501-600/956.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: Defects in CYLD are the cause of familial cylindromatosis (CYLD) also known as turban tumor syndrome or dermal eccrine cylindromatosis. CYLD is an autosomal dominant and highly tumor type-specific disorder. The tumors (known as cylindromas because of their characteristic microscopic architecture) are believed to arise from or recapitulate the appearance of the eccrine or apocrine cells of the skin that secrete sweat and scent respectively. Cylindromas arise predominantly in hairy parts of the body with approximately 90% on the head and neck. The development of a confluent mass which may ulcerate or become infected has led to the designation "turban tumor syndrome". The skin tumors show differentiation in the direction of hair structures, hence the synonym trichoepithelioma. CYLD has deubiquitinating activity.

Applications: **WB** (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (2ug/Test)**Reactivity:** Human, Mouse, Rat
(predicted: Cow, Chicken, Horse)**Predicted MW.:** 105 kDa**Subcellular Location:** Cell membrane ,Cytoplasm**— VALIDATION IMAGES —**


Sample: Cerebrum (Mouse) Lysate at 40 ug
Primary: Anti-CYLD (bs-2756R) at 1/500 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 105 kD
Observed band size: 105 kD



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (CYLD) Polyclonal Antibody, Unconjugated (bs-2756R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by microwave in sodium citrate buffer (pH6.0) ; Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes; Blocking buffer (3% BSA) at RT for 30min; Antibody incubation with (CYLD) Polyclonal Antibody, Unconjugated (bs-2756R) at 1:400 overnight at 4°C, followed by conjugation to the secondary antibody (labeled with HRP) and DAB staining.



Blank control: A431. Primary Antibody (green line): Rabbit Anti-CYLD antibody (bs-2756R)

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Dilution: 2µg /10⁶ cells; Isotype Control
Antibody (orange line): Rabbit IgG . Secondary
Antibody : Goat anti-rabbit IgG-PE Dilution: 1µg
/test. Protocol The cells were fixed with 4% PFA
(10min at room temperature)and then
permeabilized with 0.1% PBST for 20 min at
room temperature.The cells were then
incubated in 5%BSA to block non-specific
protein-protein interactions for 30 min at room
temperature .Cells stained with Primary
Antibody for 30 min at room temperature. The
secondary antibody used for 40 min at room
temperature. Acquisition of 20,000 events was
performed.