bs-0229R

[Primary Antibody]

FGF1 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 2246 SWISS: P05230

Target: FGF1

Immunogen: KLH conjugated synthetic peptide derived from human FGF1:

16-80/155.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: FGF1 also designated Heparin-binding growth factor 1 precursor (HBGF-1); Acidic fibroblast growth factor (aFGF), heparin-binding growth factors are angiogenic agents in vivo and are potent mitogens for a variety of cell types in vitro. There are differences in the tissue distribution and concentration of these 2 growth factors. Subunit si monomer.FGF1 belongs to the heparin-binding growth factors family. may play a role in neurite outgrowth; may regulate cell differentiation in the nervous system; may act in synergy with fibronectin to enhance neuronal cell adhesion.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (1ug/Test) ICC/IF (1:100-500)


Reactivity: Human, Mouse

(predicted: Rat, Sheep)


Predicted MW.: 16 kDa

Subcellular Secreted ,Cell membrane Location: ,Cytoplasm ,Nucleus


VALIDATION IMAGES




Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FGF1) Polyclonal Antibody, Unconjugated (bs-0229R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: A431. Primary Antibody (green line): Rabbit Anti-FGF1 antibody(bs-0229R) Dilution:1ug/Test; Secondary Antibody: Goat anti-rabbit IgG-FITC Dilution: 0.5ug/Test. Protocol The cells were fixed with 4% PFA



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FGF1) Polyclonal Antibody, Unconjugated (bs-0229R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



SH-SY5Y cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (FGF1) polyclonal Antibody, Unconjugated (bs-0229R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes. DAPI (blue, C02-04002) was used to stain the cell nuclei.

(10min at room temperature) and then permeabilized with 0.1% PBST for 20 min at room temperature. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

- [IF=3.8] Zhou Lijie. et al. Cellular senescence and metabolic reprogramming model based on bulk/single-cell RNA sequencing reveals PTGER4 as a therapeutic target for ccRCC. BMC CANCER. 2024 Dec;24(1):1-23 IHC; Human. 38605343
- [IF=3.8] Fengchun Liao. et al. Characterization of the Angiogenic and Proteomic Features of Circulating Exosomes in a Canine Mandibular Model of Distraction Osteogenesis. J PROTEOME RES. 2024;XXXX(XXX):XXX-XXX WB; Human. 39417529