bs-6983R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

WT1 Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 7490 **SWISS:** P19544

Target: WT1

Immunogen: KLH conjugated synthetic peptide derived from human WT1:

301-400/449.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Transcription factor that plays an important role in cellular development and cell survival. Regulates the expression of numerous target genes, including EPO. Plays an essential role for development of the urogenital system. Recognizes and binds to the DNA sequence 5'-CGCCCCGC-3'. It has a tumor suppressor as well as an oncogenic role in tumor formation. Function may be isoform-specific: isoforms lacking the KTS motif may act as transcription factors. Isoforms containing the KTS motif may bind mRNA and play a role in mRNA metabolism or splicing. Isoform 1 has lower affinity for DNA, and can bind RNA.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) **IF** (1:100-500) Flow-Cyt (1ug/test)

Reactivity: Human, Mouse

(predicted: Rat, Pig, Sheep,


Cow, Chicken, Dog)

Predicted MW.: 55 kDa

Subcellular

Location: Cytoplasm ,Nucleus


VALIDATION IMAGES




Sample: Lane 1: Mouse Testis tissue lysates Lane 2: Mouse Ovary tissue lysates Lane 3: Mouse Liver tissue lysates Lane 4: Human HL-60 cell lysates Primary: Anti-WT1 (bs-6983R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 55 kDa Observed band size: 50 kDa




Sample: TM4 Cell (Mouse) Lysate at 40 ug Primary: Anti-Wilms Tumor Protein (bs-6983R) at 1/300 dilution Secondary: IRDve800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted hand size: 55 kD Observed hand size: 48 kD




Tissue/cell: human kidney tissue: 4% Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-WT-1/Wilms Tumor Protein Polyclonal Antibody, Unconjugated(bs-6983R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (Mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block



Blank control:Molt-4. Primary Antibody (green line): Rabbit Anti-Wilms Tumor Protein antibody (bs-6983R) Dilution: 1µg/10^6 cells; Isotype

endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Wilms Tumor Protein)
Polyclonal Antibody, Unconjugated (bs-6983R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Wilms Tumor Protein)
Polyclonal Antibody, Unconjugated (bs-6983R) at 1:400 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit IgG antibody (bs-0295G-FITC) for 90 minutes, and DAPI for nuclei staining.

Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: $1\mu g$ /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at-20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at at room temperature . Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

- [IF=2.74] Rui Feng. et al. The ameliorative effect of melatonin on LPS-induced Sertoli cells inflammatory and tight junctions damage via suppression of the TLR4/MyD88/NF-kB signaling pathway in newborn calf. Theriogenology. 2022 Feb;179:103 IF; Bovine. 34871925
- [IF=2.33] Xiao, Tangli, et al. "Rapamycin promotes podocyte autophagy and ameliorates renal injury in diabetic mice." Molecular and Cellular Biochemistry (2014): 1-10. IHC; Mouse. 24850187
- [IF=2.299] Wang X et al. Influence of Wilms' tumor suppressor gene WT1 on bovine Sertoli cells polarity and tight junctions via non-canonical WNT signaling pathway. Theriogenology. 2019 Jul 8;138:84-93. ICC; COW. 31302435
- [IF=2.299] Wang X et al. 3, 3', 5-Triiodo-L-thyronine affects polarity proteins of bovine Sertoli cells via WT1/non-canonical Wnt signaling pathway. Theriogenology. 2020 Feb 21;148:8-17. WB; Bovine. 32126395
- [IF=1.723] Wang X et al. Wilms' tumour 1 (WT1) negatively regulates the expression of connexin 43 via a non-canonical Wnt signalling pathway in cultured bovine Sertoli cells. Reprod Fertil Dev. 2020 Feb 6. WB,ICC; Bovine. 32023428