



SMAD1 + SMAD5 Rabbit pAb

Catalog Number: bs-2973R
Target Protein: SMAD1 + SMAD5
Concentration: 1mg/ml
Form: Liquid
Host: Rabbit
Clonality: Polyclonal
Isotype: IgG
Applications: WB (1:200-1000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500), ICC/IF (1:100)
Reactivity: Human, Mouse, Rabbit (predicted:Rat, Cow, Chicken, Dog, Horse)
Predicted MW: 51 kDa
Entrez Gene: 4090
Swiss Prot: Q99717
Source: KLH conjugated synthetic peptide derived from human SMAD5: 361-465/465.
Purification: affinity purified by Protein A
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Background: The protein encoded by this gene is involved in the transforming growth factor beta signaling pathway that results in an inhibition of the proliferation of hematopoietic progenitor cells. The encoded protein is activated by bone morphogenetic proteins type 1 receptor kinase, and may be involved in cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014]


VALIDATION IMAGES



Sample: Lane 1: Mouse Cerebrum tissue lysates Lane 2: Rat Cerebrum tissue lysates Primary: Anti-SMAD1 + SMAD5 (bs-2973R) at 1/200 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 51 kDa Observed band size: 55 kDa



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SMAD1 + SMAD5) Polyclonal Antibody, Unconjugated (bs-2973R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (SMAD1 + SMAD5) polyclonal Antibody, Unconjugated (bs-2973R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

PRODUCT SPECIFIC PUBLICATIONS

[IF=2.9] Zhen Liu. et al. Enhanced bacteriostasis and osseointegrative properties of SiRNA-modified polyetheretherketone surface for implant applications. PLOS ONE. 2024 Dec;19(12):e0314091 WB ; Rat . 39636795

[IF=1.663] Yu M et al. BMP4 mutations in tooth agenesis and low bone mass. Arch Oral Biol. 2019 Jul;103:40-46. WB ; Human . 31128441