

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Shh Rabbit pAb

Catalog Number: bs-1544R

Target Protein: Shh

Concentration: 1mg/ml

Form: Liquid Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat (predicted:Rabbit, Cow, Chicken, Horse)

Predicted MW: 19/47 kDa

Entrez Gene: 6469

Swiss Prot: Q15465

Source: KLH conjugated synthetic peptide derived from human Shh: 21-120/462.

Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: SHH binds to the patched (PTC) receptor, which functions in association with smoothened


(SMO), to activate the transcription of target genes. In the absence of SHH, PTC represses the constitutive signaling activity of SMO. Also regulates another target, the gli oncogene. Intercellular signal essential for a variety of patterning events during development: signal produced by the notochord that induces ventral cell fate in the neural tube and somites, and the polarizing signal for patterning of the anterior-posterior axis of the developing limb bud. Displays both floor plate- and motor neuron-inducing activity. The threshold concentration of N-product required for motor neuron induction is 5-fold lower than that required for floor

plate induction (By similarity).


VALIDATION IMAGES




Sample: Embryo(Mouse) tissue at 30ug; Primary: Anti-Shh (bs-1544R) at 1:300 dilution; Secondary: HRP conjugated Goat Anti-Rabbit IgG(bs-0295G-HRP) at 1:5000 dilution; Predicted band size: 19/47kD Observed band size: 50kD



Sample:Stomach (Mouse) Lysate at 40 ug Primary: Anti-Shh(bs-1544R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 19/47kD Observed band size: 49kD



Paraformaldehyde-fixed, paraffin embedded (Mouse stomach); Antigen retrieval by microwave in sodium citrate buffer (pH6.0); Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes; Blocking buffer (3% BSA) at RT for 30min; Antibody incubation with (Shh) Polyclonal Antibody, Unconjugated (bs-1544R) at 1:400 overnight at 4°C, followed by conjugation to the secondary antibody (labeled with HRP) and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Shh) Polyclonal Antibody, Unconjugated (bs-1544R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

PRODUCT SPECIFIC PUBLICATIONS

[IF=8.98] Weiss, Robert M., et al. "Brg1 Determines Urothelial Cell Fate during Ureter Development." Journal of the American Society of Nephrology: JASN (2013). Other; ="" . 23449535

[IF=8.718] Chao Sun. et al. WISP-1 induced by mechanical stress contributes to fibrosis and hypertrophy of the ligamentum flavum through Hedgehog-Gli1 signaling. Exp Mol Med. 2021 Jun;53(6):1068-1079 WB; Human. 34158608

[IF=5.682] Qi, Yan. et al. Neuroprotective Effect of Sonic Hedgehog Mediated PI3K/AKT Pathway in Amyotrophic Lateral Sclerosis Model Mice. MOL NEUROBIOL. 2022 Sep;:1-12 WB; MOUSE . 36056982

[IF=4.56] Wang, H., et al. "Baicalin Attenuates Alcoholic Liver Injury through Modulation of Hepatic Oxidative Stress, Inflammation and Sonic Hedgehog Pathway in Rats." Cellular Physiology and Biochemistry 39.3 (2016): 1129-1140. WB; = "Rat". 27576501

[IF=4.46] Du, Wen - Zhong, et al. "Curcumin Suppresses Malignant Glioma Cells Growth and Induces Apoptosis by Inhibition of SHH/GLI1

Signaling Pathway in Vitro and Vivo." CNS Neuroscience & Therapeutics (2013). WB; = "Human" . 24165291