bs-2758R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Cdc25A Rabbit pAb

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 993 **SWISS:** P30304

Target: Cdc25A

Immunogen: KLH conjugated synthetic peptide derived from human cdc25A:

451-524/524.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: CDC25A is a member of the CDC25 family of phosphatases. CDC25A is required for progression from G1 to the S phase of the cell cycle. It activates the cyclin-dependent kinase CDC2 by removing two phosphate groups. CDC25A is specifically degraded in response to DNA damage, which prevents cells with chromosomal abnormalities from progressing through cell division. CDC25A is an

oncogene, although its exact role in oncogenesis has not been demonstrated. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Applications: WB (1:500-2000)

IHC-P (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500) Flow-Cyt (2ug/Test)

Reactivity: Human, Mouse, Rat


(predicted: Rabbit, Cow,

Chicken, Dog)


Predicted MW.: 59 kDa

Subcellular Location: Cytoplasm ,Nucleus


VALIDATION IMAGES -




Sample: Testis (Mouse) Lysate at 40 ug Primary: Anti- Cdc25A (bs-2758R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 59 kD Observed band size: 61 kD



Paraformaldehyde-fixed, paraffin embedded (Rat small intestine): Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Cdc25A) Polyclonal Antibody, Unconjugated (bs-2758R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Cdc25A) Polyclonal Antibody, Unconjugated (bs-2758R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control (black line) :HepG2. Primary Antibody (green line): Rabbit Anti-Cdc25A antibody (bs-2758R) Dilution:2ug/Test; Secondary Antibody (white blue line): Goat anti-rabbit IgG-AF488 Dilution: 0.5ug/Test.

Isotype control (orange line): Normal Rabbit IgG Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

- SELECTED CITATIONS -

- [IF=11.205] Cong Lan. et al. Inhibition of DYRK1A, via histone modification, promotes cardiomyocyte cell cycle activation and cardiac repair after myocardial infarction. EBIOMEDICINE. 2022 Aug;82:104139 WB;Rat. 35810562
- [IF=3.98] Wei, Jialiu, et al. "Endosulfan induces cell dysfunction through cycle arrest resulting from DNA damage and DNA damage response signaling pathways." Science of The Total Environment 589 (2017): 97-106. WB; Human. 28273598