

bs-43649R**[Primary Antibody]****TNFSF13B Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: Flow-Cyt (1ug/Test) ICC/IF (1:50-200) ELISA (1:5000-10000) Reactivity: Human Predicted MW.: 34 kDa Subcellular Location: Secreted ,Cell membrane
Clonality: Polyclonal		
GeneID: 10673	SWISS: Q9Y275	
Target: TNFSF13B		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS(pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.		
Background: The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for receptors TNFRSF13B/TACI, TNFRSF17/BCMA, and TNFRSF13C/BAFFR. This cytokine is expressed in B cell lineage cells, and acts as a potent B cell activator. It has been also shown to play an important role in the proliferation and differentiation of B cells. [provided by RefSeq].		

— VALIDATION IMAGES —

4% Paraformaldehyde-fixed U937 (H) cell;
Antibody incubation with (TNFSF13B) polyclonal
Antibody, unconjugated (bs-43649R) 1:100, 90
min at 37°C; followed by conjugated Goat Anti-
Rabbit IgG antibody (green, bs-40295G-FITC) at
37°C for 90 min, DAPI (blue, C02-04002) was used
to stain the cell nuclei. PBS instead of the
primary antibody was used as the blank control.