

bsm-61530R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

CEBP beta Recombinant Rabbit mAb**— DATASHEET —**


Host: Rabbit
Clonality: Recombinant
GeneID: 1051
Target: CEBP beta
Immunogen: A synthesized peptide derived from human CEBP beta: 310-345.
Purification: affinity purified by Protein A
Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles.
Background: Important transcription factor regulating the expression of genes involved in immune and inflammatory responses.

Isotype: IgG**CloneNo.:** 16G3**SWISS:** P17676


Applications: WB (1:500-2000)
Flow-Cyt (1:50-100)
ICC/IF (1:50-200)
IP (1:20-50)

Reactivity: Human**Predicted
MW.:** 36**Subcellular
Location:** Nucleus**— VALIDATION IMAGES —**

25 ug total protein per lane of various lysates (see on figure) probed with CEBP beta monoclonal antibody, unconjugated (bsm-61530R) at 1:2000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



4% Paraformaldehyde-fixed MCF-7 (H) cell; Triton X-100 at r.t. for 20 min; Antibody incubation with (CEBP beta) monoclonal Antibody, unconjugated (bsm-61530R) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, bs-60295G-BF488) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.



The MCF-7 (H) cells were fixed with 4% PFA (10 min at r.t.) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C, the cells then were incubated in 5%BSA to block non-specific protein-protein interactions (30 min at r.t.). Primary Antibody (green): Rabbit Anti-CEBP beta antibody (bsm-61530R): 1 µg/10⁶ cells; Secondary Antibody (white blue): Goat anti-Rabbit IgG-BF488 (bs-60295G-BF488): 1 µg/test. Isotype Control (orange): Rabbit IgG (bs-0295P). Blank control (black): PBS. Acquisition of 20,000 events was performed.