

bs-1119R

[Primary Antibody]

PAF Rabbit pAb

BioSS
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Target: PAF

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Platelet-activating factor, a compound that reduces inflammation by increasing permeability of blood vessels and contracting various involuntary muscles such as those in airways. It is a phospholipid with a wide range of proinflammatory actions, is immediately degraded and inactivated in vivo by PAF acetylhydrolase.

Applications: IHC-P (1:100-500)


IHC-F (1:100-500)

IF (1:100-500)

Reactivity: Rat (predicted: PAF)

Predicted MW.: 0.524 kDa

— VALIDATION IMAGES —



Tissue/cell: rat colon tissue;4%
Paraformaldehyde-fixed and paraffin-embedded
Antigen retrieval: citrate buffer (0.01M, pH 6.0),
Boiling bathing for 15min; Block endogenous
peroxidase by 3% Hydrogen peroxide for 30min;
Blocking buffer (normal goat serum) at 37°C for
20 min Incubation: Anti-PAF Polyclonal
Antibody, Unconjugated(bs-1119R) 1:200,
overnight at 4°C, followed by conjugation to the
secondary antibody and DAB staining