

bs-0860R

[Primary Antibody]

HHV8 ORF50 Rabbit pAb



www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

<p>Host: Rabbit</p> <p>Clonality: Polyclonal</p> <p>Target: HHV8 ORF50</p> <p>Immunogen: KLH conjugated synthetic peptide derived from HHV8/ORF50: 601-691/691.</p> <p>Purification: affinity purified by Protein A</p> <p>Concentration: 1mg/ml</p> <p>Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.</p> <p>Background: HHV8 has been found to be associated with three different diseases observed in AIDS patients; kaposi's sarcoma, primary effusion lymphoma (which is a rare type of non-Hodgkin lymphoma affecting the body cavities) and multicentric Castleman's disease. To date there is much evidence to support a direct role for HHV8 in kaposi's sarcoma.</p>	<p>Isotype: IgG</p> <p>Applications: ELISA (1:5000-10000)</p> <p>Reactivity: (predicted: herpesvirus 8type P)</p> <p>Predicted MW.: 74 kDa</p>
--	---

— SELECTED CITATIONS —

- **[IF=5.64]** Su-Kyung Kang. et al. Dimethyl Sulfoxide Enhances Kaposi' s Sarcoma-Associated Herpesvirus Production During Lytic Replication. Front Microbiol. 2021; 12: 778525 WB ;Human. 34975802