bsm-61767R

[Primary Antibody]

Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

TLE1 Recombinant Rabbit mAb

- DATASHEET -

Host: Rabbit Isotype: IgG
Clonality: Recombinant CloneNo.: 7D8
GeneID: 7088 SWISS: Q04724

Target: TLE1

Immunogen: A synthesized peptide derived from human TLE1: 200-218.

Purification: affinity purified by Protein A

Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium

chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid

repeated freeze/thaw cycles.

Background: Transcriptional corepressor that binds to a number of transcription

factors. Inhibits NF-kappa-B-regulated gene expression. Inhibits the transcriptional activation mediated by FOXA2, and by CTNNB1 and

TCF family members in Wnt signaling.

Applications: WB (1:500-2000)


IHC-P (1:50-200) IHC-F (1:50-200) IF (1:50-200) ICC/IF (1:50-200)

Reactivity: Human


Predicted 83

Subcellular Location: Nucleus


VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with TLE1 monoclonal antibody, unconjugated (bsm-61767R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Endometrial Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with TLE1 Monoclonal Antibody, Unconjugated(bsm-61767R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Glioma; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with TLE1 Monoclonal Antibody, Unconjugated(bsm-61767R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.