bsm-61851R

[Primary Antibody]

BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Galectin 7 Recombinant Rabbit mAb

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Recombinant

GenelD: 3963 **SWISS:** P47929

Target: Galectin 7

Immunogen: A synthesized peptide derived from human Galectin 7: 1-50/136.

Purification: affinity purified by Protein A

Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium

chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid

repeated freeze/thaw cycles.

Background: Could be involved in cell-cell and/or cell-matrix interactions

necessary for normal growth control. Pro-apoptotic protein that functions intracellularly upstream of JNK activation and cytochrome

c release.

Applications: WB (1:500-2000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:50-200) IP (1:20-50)


Reactivity: Human

Predicted MW.: 15

Subcellular Secreted ,Cytoplasm

Location: ,Nucleus


VALIDATION IMAGES -



Western blot analysis of HACAT cell lysate. Using Galectin 7 (bsm-61851R) monoclonal antibody at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Tonsil; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Galectin 7 Monoclonal Antibody, Unconjugated (bsm-61851R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Cervical Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with Galectin 7 Monoclonal Antibody, Unconjugated (bsm-61851R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.