


**bsm-62408R****[ Primary Antibody ]****BioSS**  
**ANTIBODIES**

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

**Formyl-Histone H2B type 2E (Lys117)  
Recombinant Rabbit mAb****— DATASHEET —****Host:** Rabbit**Isotype:** IgG**Clonality:** Recombinant**GeneID:** 8349**SWISS:** Q16778**Target:** Formyl-Histone H2B type 2E (Lys117)**Immunogen:** A synthesized peptide derived from human Histone H2B type 2E around the formylation site of K117: GT-K(Fo)-AVT.**Purification:** affinity purified by Protein A**Storage:** 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles.**Background:** Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.**Applications:** **WB** (1:1000-2000)  
**ICC/IF** (1:50-200)**Reactivity:** Human, Mouse**Predicted MW.:** 14**— VALIDATION IMAGES —**

Western blot analysis of HeLa cell lysate. Using Formyl-Histone H2B type 2E (Lys117) (bsm-62408R) monoclonal antibody at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.