bs-11518R

[Primary Antibody]

www.bioss.com.cn sales@bioss.com.cn

techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

FOXD3 Rabbit pAb

GenelD: 27022 SWISS: Q9UJU5

Target: FOXD3

Immunogen: KLH conjugated synthetic peptide derived from human FOXD3:

151-230/478.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Embryonic stem cells require the forkhead transcriptional

regulator FoxD3 for survival. Following gastrulation, FoxD3 generally gets downregulated, except in the neural crest. A variety of growth factors induce FoxD3 expression, including FGF8 and SNAIL, maintaining the effected cells in an undifferentiated state. Thus defects in FoxD3 induction may cause premature

differentiation and/or migration-asociated birth defects.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500) **IF** (1:100-500) ICC/IF (1:25)

Reactivity: Human, Mouse, Rat

(predicted: Rabbit, Cow,

Chicken, Dog)


Predicted 48 kDa

Subcellular Nucleus Location:


VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (FOXD3) Polyclonal Antibody, Unconjugated (bs-11518R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (FOXD3) Polyclonal Antibody, Unconjugated (bs-11518R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Hela cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (FOXD3) polyclonal Antibody, Unconjugated (bs-11518R) 1:25, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.

— SELECTED CITATIONS ———

• [IF=2.688] Wang, et al. FOXD3/FOXD4 is required for the development of hindgut in the rat model of anorectal malformation.(2018) Experimental Biology and Medicine. 243:327-333. IHC; Rat. 29307282