bs-11522R

[Primary Antibody]

www.bioss.com.cn

NPY4R Rabbit pAb

sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 100996758 **SWISS:** P50391

Target: NPY4R

Immunogen: KLH conjugated synthetic peptide derived from human NPY4R:

151-250/375. < Extracellular >

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Pancreatic polypeptide (PP), neuropeptide Y (NPY), and peptide YY

(PYY) are related 36-amino acid hormones. A number of structurally related receptors for these peptides have been isolated, NPY1-R, NPY2-R, NPY3-R, NPY4-R, NPY5-R, and NPY6-R. NPY4-R is expressed in several human tissues, including brain, coronary artery, and ileum. NPY4-R maps to human chromosome 10g11.2. NPY-5R, isolated from rat hypothalamus, encodes a 456amino acid protein with less than 35% overall identity to known Ytype receptors. The human NPY5-R sequence is nearly identical to, but in the opposite orientation from, that of the human NPY1-R sequence. NPY5-R localizes to the paraventricular hypothalamic nucleus, the lateral hypothalamus, and other locations consistent with a role in the control of feeding behavior. The gene which encodes NPY5-R maps to human chromosome 4q32.2. NPY6-R is abundantly expressed in human heart and skeletal muscle and the gene which encodes NPY6-R maps to human chromosome 5q31.

Applications: WB (1:500-2000)


Reactivity: Mouse (predicted: Human,

Rabbit)

Predicted 42 kDa MW.:

Subcellular Location: Cell membrane

VALIDATION IMAGES -



Sample: Small intestine (Mouse) Lysate at 40 ug Large intestine (Mouse) Lysate at 40 ug Primary: Anti- NPY4R (bs-11522R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 42 kD Observed band size: 62 kD