


bsm-62477R**[Primary Antibody]****CD133 Recombinant Rabbit mAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

DATASHEET**Host:** Rabbit**Isotype:** IgG**Clonality:** Recombinant**CloneNo.:** 11G1**GeneID:** 8842**SWISS:** O43490**Target:** CD133**Immunogen:** A synthesized peptide derived from human CD133: 250-380.**Purification:** affinity purified by Protein A**Storage:** 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles.**Background:** May play a role in cell differentiation, proliferation and apoptosis. Binds cholesterol in cholesterol-containing plasma membrane microdomains and may play a role in the organization of the apical plasma membrane in epithelial cells. During early retinal development acts as a key regulator of disk morphogenesis. Involved in regulation of MAPK and Akt signaling pathways. In neuroblastoma cells suppresses cell differentiation such as neurite outgrowth in a RET-dependent manner.**Applications:** **WB** (1:1000-5000)
IHC-P (1:200-1000)
IHC-F (1:200-1000)
IF (1:200-1000)**Reactivity:** Human**Predicted
MW.:** 97**Subcellular
Location:** Cell membrane**VALIDATION IMAGES**

25 ug total protein per lane of various lysates (see on figure) probed with CD133 monoclonal antibody, unconjugated (bsm-62477R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Pancreas; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD133 Monoclonal Antibody, Unconjugated (bsm-62477R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with CD133 Monoclonal Antibody, Unconjugated (bsm-62477R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.