bsm-62700R

[Primary Antibody]

TRIP1 Recombinant Rabbit mAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET -Host: Rabbit Isotype: IgG Applications: WB (1:1000-1:2000) **IHC-P** (1:100-1:200) Clonality: Recombinant IHC-F (1:100-1:200) GenelD: 5705 SWISS: P62195 **IF** (1:50-1:200) Flow-Cyt (1:20-1:100) Target: TRIP1 ICC/IF (1:50-1:200) Immunogen: A synthesized peptide derived from human PSMC5: 1-26. IP (1:20-1:50) Purification: affinity purified by Protein A Reactivity: Human, Mouse, Rat Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles. Predicted MW.: 46 Background: The 26S protease is involved in the ATP-dependent degradation of ubiquitinated proteins. The regulatory (or ATPase) complex confers ATP dependency and substrate specificity to the 26S complex. Subcellular Location: Cytoplasm ,Nucleus

- VALIDATION IMAGES -



Western blot analysis of A431 cell lysate. Using TRIP1 (bsm-62700R) monoclonal antibody at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.