bsm-62910R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

CD10 Recombinant Rabbit mAb

- DATASHEET -

Host: Rabbit Isotype: IgG
Clonality: Recombinant CloneNo.: 6E16
GeneID: 4311 SWISS: P08473

Target: CD10

Immunogen: A synthesized peptide derived from human Neprilysin: 50-500/750.

Purification: affinity purified by Protein A

Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid

repeated freeze/thaw cycles.

Background: Thermolysin-like specificity, but is almost confined on acting on

polypeptides of up to 30 amino acids. Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by

cleavage of a Gly-Phe bond.

Applications: WB (1:500-2000)


IHC-P (1:50-200) IHC-F (1:50-200) IF (1:50-200) IP (1:20-50)

Reactivity: Human


Predicted 86

Subcellular Location: Cell membrane


VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with CD10 monoclonal antibody, unconjugated (bsm-62910R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Small Intestine; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD10 Monoclonal Antibody, Unconjugated(bsm-62910R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Placenta; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with CD10 Monoclonal Antibody, Unconjugated(bsm-62910R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.