bsm-62975R

[Primary Antibody]

BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

EDG3 Recombinant Rabbit mAb

- DATASHEET -

Host: Rabbit Isotype: IgG
Clonality: Recombinant CloneNo.: 9E8
GeneID: 1903 SWISS: Q99500

Target: EDG3

Immunogen: A synthesized peptide derived from human EDG3: 330-378/378.

Purification: affinity purified by Protein A

Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium

chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid

repeated freeze/thaw cycles.

Background: Receptor for the lysosphingolipid sphingosine 1-phosphate (S1P).

S1P is a bioactive lysophospholipid that elicits diverse physiological effect on most types of cells and tissues. When expressed in rat HTC4

proliferation and suppression of apoptosis.

hepatoma cells, is capable of mediating S1P-induced cell

Applications: WB (1:1000-1:2000)


IHC-P (1:100-1:200) IHC-F (1:100-1:200) IF (1:100-1:200) ICC/IF (1:50-1:200)

Reactivity: Human


Predicted MW.: 42

Subcellular Location: Cell membrane


- VALIDATION IMAGES -



Western blot analysis of HepG2 cell lysate. Using EDG3 (bsm-62975R) monoclonal antibody at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Heart; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with EDG3 Monoclonal Antibody, Unconjugated(bsm-62975R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with EDG3 Monoclonal Antibody, Unconjugated(bsm-62975R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.