bsm-62978R

[Primary Antibody]

Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

FATP4 Recombinant Rabbit mAb

- DATASHEET -

Host: Rabbit Isotype: IgG
Clonality: Recombinant CloneNo.: 20C7
GeneID: 10999 SWISS: Q6P1M0

Target: FATP4

Immunogen: A synthesized peptide derived from human FATP 4: 400-643.

Purification: affinity purified by Protein A

Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium

chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid

repeated freeze/thaw cycles.

Background: Mediates the levels of long-chain fatty acids (LCFA) in the cell by

facilitating their transport across cell membranes. Appears to be the principal fatty acid transporter in small intestinal enterocytes. Also functions as an acyl-CoA ligase catalyzing the ATP-dependent formation of fatty acyl-CoA using LCFA and very-long-chain fatty acids (VLCFA) as substrates, which prevents fatty acid efflux from cells

and might drive more fatty acid uptake.

Applications: WB (1:500-2000)

Flow-Cyt (1:50-100) ICC/IF (1:50-200)


Reactivity: Mouse (predicted: Human,

Rat)

Predicted 72

Subcellular Location: Cell membrane

VALIDATION IMAGES



25 ug total protein per lane of various lysates (see on figure) probed with FATP4 monoclonal antibody, unconjugated (bsm-62978R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.