[Primary Antibody]





www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

– DATASHEET –––––		400-901-9800
Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal	C C	ICC/IF (1:100-500) ELISA (1:5000-10000)
GenelD: 2693	SWISS: Q6ISR8	
Target: Ghrelin Receptor		Reactivity: (predicted: Human, Mouse, Rat, Rabbit, Pig, Sheep,
Immunogen: KLH conjugated synthetic peptide derived from human Ghrelin Receptor: 75-135/366. < Extracellular >		Cow, Horse)
Purification: affinity purified by Protein A		Predicted MW.: ⁴¹ kDa Subcellular Location: Cell membrane
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: This gene encodes a member of the G-protein coupledreceptor family. The encoded protein may play a role in energyhomeostasis and regulation of body weight. Two identifiedtranscript variants are expressed in several tissues and areevolutionary conserved in fish and swine. One transcript, 1a, excises an intron and encodes the functional protein; this proteinis the receptor for the Ghrelin ligand and defines a neuroendocrinepathway for growth hormone release. The second transcript (1b)retains the intron and does not function as a receptor for Ghrelin; however, it may function to attenuate activity of isoform 1a.Mutations in this gene are associated with autosomal idiopathicshort stature.[provided by RefSeq, Apr 2010].		

- SELECTED CITATIONS -

- [IF=5.81] Shengyan Xi. et al. The Ciji-Hua' ai-Baosheng II Formula Attenuates Chemotherapy-Induced Anorexia in Mice With H22 Hepatocellular Carcinoma. Front Pharmacol. 2021; 12: 715824 IHC ;Mouse. 34489705
- [IF=4.994] He Zhang. et al. Effect of feeding regimen on circadian activity rhythms of food anticipatory by ghrelin hormone in a pig model. 2022 Mar 05 WB ;Pig. 35249475
- [IF=4.546] Quancheng Liu. et al. FumDSB Can Reduce the Toxic Effects of Fumonisin B1 by Regulating Several Brain-Gut Peptides in Both the Hypothalamus and Jejunum of Growing Pigs. Toxins. 2021 Dec;13(12):874 WB,IHC ;Pig. 34941712
- **[IF=2.784]** Liu et al. Ghrelin protects the myocardium with hypoxia/reoxygenation treatment through upregulating the expression of growth hormone, growth hormone secretagogue receptor and insulin-like growth factor-1, and promoting the phosphorylation of protein kinase B. (2018) Int.J.Mol.Med. 42:3037-3046 WB,IHC ;Rat. 30272367
- [IF=3] He Zhang. et al. Inhibition of ghrelin activity by the receptor antagonist [D-Lys3]-GHRP-6 enhances hepatic fatty acid oxidation and gluconeogenesis in a growing pig model. PEPTIDES. 2023 Aug;166:171041 WB ;Pig. 37301480