bsm-62979R

## [ Primary Antibody ]


## FATP4 Recombinant Rabbit mAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

- DATASHEET		400-901-9800
Host: Rabbit	<b>lsotype:</b> lgG	Applications: WB (1:1000-2000)
Clonality: Recombinant		IHC-P (1:100-200) IHC-F (1:100-200)
GenelD: 10999	SWISS: Q6P1M0	<b>IF</b> (1:50-200)
Target: FATP4		ICC/IF (1:50-200) IP (1:20-50)
Immunogen: A synthesized peptide derived from human FATP 4: 400-643. Purification: affinity purified by Protein A		Reactivity: Human
<b>Storage:</b> 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles.		Predicted MW.: <sup>72</sup>
<b>Background:</b> Mediates the levels of long-chain fatty acids (LCFA) in the cell by facilitating their transport across cell membranes. Appears to be the principal fatty acid transporter in small intestinal enterocytes. Also functions as an acyl-CoA ligase catalyzing the ATP-dependent formation of fatty acyl-CoA using LCFA and very-long-chain fatty acids (VLCFA) as substrates, which prevents fatty acid efflux from cells and might drive more fatty acid uptake.		Subcellular Location: <sup>Cell</sup> membrane

## - VALIDATION IMAGES -



Western blot analysis of HepG2 cell lysate. Using FATP4 (bsm-62979R) monoclonal antibody at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.