

bsm-62996R

[Primary Antibody]

TRAF2 Recombinant Rabbit mAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

Host: Rabbit

Clonality: Recombinant

GeneID: 7186

Target: TRAF2

Immunogen: A synthesized peptide derived from human TRAF2: 1-28.

Purification: affinity purified by Protein A

Storage: 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid repeated freeze/thaw cycles.

Background: Regulates activation of NF-kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis.

Isotype: IgG

CloneNo.: 15A5

SWISS: Q12933

Applications: WB (1:500-2000)

IHC-P (1:50-200)

IHC-F (1:50-200)

IF (1:50-200)

Flow-Cyt (1:20-50)

ICC/IF (1:50-200)


IP (1:20-50)

Reactivity: Human, Mouse, Rat


**Predicted
MW.:** 56

**Subcellular
Location:** Cytoplasm


— VALIDATION IMAGES —



25 ug total protein per lane of various lysates (see on figure) probed with TRAF2 monoclonal antibody, unconjugated (bsm-62996R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with TRAF2 Monoclonal Antibody, Unconjugated (bsm-62996R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with TRAF2 Monoclonal Antibody, Unconjugated (bsm-62996R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.