## bsm-63004R

## [ Primary Antibody ]

# Bioss ANTIBODIES

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

### **UBA1** Recombinant Rabbit mAb

- DATASHEET -

Host: Rabbit Isotype: IgG
Clonality: Recombinant CloneNo.: 7D16
GeneID: 7317 SWISS: P22314

Target: UBA1

Immunogen: A synthesized peptide derived from human UBA1: 200-255/1058.

Purification: affinity purified by Protein A

**Storage:** 10mM phosphate buffered saline(pH 7.4) with 150mM sodium chloride, 0.05% BSA, 0.02% Proclin300 and 50% glycerol. Store at 4°C for short term. Store at -20°C for long term. Avoid

repeated freeze/thaw cycles.

**Background:** Activates ubiquitin by first adenylating its C-terminal glycine residue

with ATP, and thereafter linking this residue to the side chain of a cysteine residue in E1, yielding an ubiquitin-E1 thioester and free

AMP.

**Applications: WB** (1:500-2000)


IHC-P (1:50-200) IHC-F (1:50-200) IF (1:50-200) Flow-Cyt (1:50-100) ICC/IF (1:50-200)

Reactivity: Human, Mouse, Rat

Predicted MW.: 118

Subcellular Cytoplasm ,Nucleus


#### VALIDATION IMAGES -



25 ug total protein per lane of various lysates (see on figure) probed with UBA1 monoclonal antibody, unconjugated (bsm-63004R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human ovarian cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with UBA1 Monoclonal Antibody, Unconjugated(bsm-63004R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Kidney; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with UBA1 Monoclonal Antibody, Unconjugated(bsm-63004R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.