

bs-17488R**[Primary Antibody]****DC-SIGNR1/CD209b Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		
GeneID: 69165	SWISS: Q8CJ91	
Target: DC-SIGNR1/CD209b		
Immunogen: KLH conjugated synthetic peptide derived from mouse CD209b: 51-150/325.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		Reactivity: Mouse
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		Predicted MW.: 37 kDa
Background: Antigen-presenting cells are localized in essentially every tissue, where they operate at the interface of innate and acquired immunity by capturing pathogens and presenting pathogen-derived peptides to T cells. Dendritic cells capture antigens or viruses in peripheral tissue and transport them to lymphoid organs, an event that induces cellular T cell responses. The mouse CD209 family of cell adhesion receptors consists of CD209a (also known as DC-SIGN), CD209b, CD209c, CD209d, CD209e, CD209f and CD209g, all of which function to mediate the endocytosis and subsequent degradation of pathogens within lysosomal compartments. There are two human CD209 proteins, designated DC-SIGN and DC-SIGNR, which function in a similar manner to the mouse proteins.		Subcellular Location: Cell membrane

— VALIDATION IMAGES —

Sample: Pancreas (Mouse) Lysate at 40 ug
Primary: Anti-DC-SIGNR1/CD209b(bs-17488R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 37kD Observed band size: 37kD



Sample: Pancreas (Mouse) Lysate at 40 ug
Primary: Anti-DC-SIGNR1/CD209b(bs-17488R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 37 kD Observed band size: 37 kD

— SELECTED CITATIONS —

- **[IF=5.6]** Xiumeng Hua. et al. STING regulates the transformation of the proinflammatory macrophage phenotype by HIF1A into autoimmune myocarditis. INT IMMUNOPHARMACOL. 2023 Aug;121:110523 IHC ;Mouse. 37354779