bsm-33019R

# [ Primary Antibody ]

# BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

# - DATASHEET -

Host: Rabbit

Clonality: Recombinant CloneNo.: 15F4

**GFP Recombinant Rabbit mAb** 

Target: GFP

Purification: affinity purified by Protein A

**Storage:** 0.01M TBS(pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20 °C for one year. Avoid repeated

freeze/thaw cycles.

**Background:** Enhanced Blue Fluorescent Protein (EBFP) emits a strong blue fluorescence. EBFP can be used as protein "tags" to study the sub-


fluorescence. EBFP can be used as protein "tags" to study the subcellular localization of proteins and/or their translocation upon stimulation or as markers for transfection in transient and stable


expression systems.

**Applications: WB** (1:1000-10000)

Reactivity: Belt Jellyfish

## VALIDATION IMAGES





20 ng rGFP-His protein (bs-33009P) per lane probed with GFP monoclonal antibody respectively, unconjugated (bsm-33019R) at 1:1000 dilution and 4°C overnight incubation. Followed by corresponding conjugated secondary antibody incubation at r.t. for 60 min.

50 ng rGFP-His protein (bs-33009P) per lane probed with GFP monoclonal antibody respectively, unconjugated (bsm-33019R) at 1:10000 dilution and 4°C overnight incubation. Followed by corresponding conjugated secondary antibody incubation at r.t. for 60 min.

### — SELECTED CITATIONS –

• [IF=5.7] Yu Xia. et al. Importance of Selenoprotein O in Regulating Hmgb1: A New Direction for Modulating ROS-Dependent NETs Formation to Aggravate the Progression of Acute Liver Inflammation. J AGR FOOD CHEM. 2025;73(15):9382–9397 WB; Mouse. 40189811