


bsm-33019R**[Primary Antibody]****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

GFP Recombinant Rabbit mAb**— DATASHEET —****Host:** Rabbit**Clonality:** Recombinant**CloneNo.:** 15F4**Target:** GFP**Purification:** affinity purified by Protein A**Storage:** 0.01M TBS(pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.**Background:** Enhanced Blue Fluorescent Protein (EBFP) emits a strong blue fluorescence. EBFP can be used as protein "tags" to study the sub-cellular localization of proteins and/or their translocation upon stimulation or as markers for transfection in transient and stable expression systems.**Applications:** **WB** (1:1000-10000)
ELISA (1:1000-5000)**Reactivity:** Species independent**Predicted MW.:** 27 kDa**— VALIDATION IMAGES —**

20 ng rGFP-His protein (bs-33009P) per lane probed with GFP monoclonal antibody respectively, unconjugated (bsm-33019R) at 1:1000 dilution and 4°C overnight incubation. Followed by corresponding conjugated secondary antibody incubation at r.t. for 60 min.



50 ng rGFP-His protein (bs-33009P) per lane probed with GFP monoclonal antibody respectively, unconjugated (bsm-33019R) at 1:10000 dilution and 4°C overnight incubation. Followed by corresponding conjugated secondary antibody incubation at r.t. for 60 min.

— SELECTED CITATIONS —

- **[IF=5.7]** Yu Xia. et al. Importance of Selenoprotein O in Regulating Hmgb1: A New Direction for Modulating ROS-Dependent NETs Formation to Aggravate the Progression of Acute Liver Inflammation. J AGR FOOD CHEM. 2025;73(15):9382-9397 WB ;Mouse. 40189811