

bsm-34186M

[Primary Antibody]

MUM1 Mouse mAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

DATASHEET

Host: Mouse

Clonality: Monoclonal

GeneID: 84939

Target: MUM1

CloneNo.: 9H6

SWISS: Q2TAK8

Purification: affinity purified by Protein G

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Munc 13s are homologues of the *C. elegans* unc-13 gene product. Three brain specific isoforms are expressed in rat: Munc 13-1, -13-2, and -13-3. Munc 13-1 is a peripheral membrane protein that is enriched in synaptosomes and localized to plasma membranes but absent from synaptic vesicles. It has multiple regulatory domains (C1 and C2 domains) that may mediate diacylglycerol, phospholipid, and calcium binding. The C terminus of Munc 13-1 interacts directly with a putative coiled coil domain in the N-terminal part of syntaxin. It has been shown that Munc 13 is involved in priming synaptic vesicles. These activated vesicles can be released immediately on Ca²⁺ influx.

Applications: IHC-P (1:100-500)

IHC-F (1:100-500)


IF (1:100-500)

Reactivity: Human


**Predicted
MW.:** 79 kDa

**Subcellular
Location:** Cell membrane ,Cytoplasm

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded Human lymph; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with MUM1 Monoclonal Antibody, Unconjugated (bsm-34186M) at 1:200 overnight at 4°C, followed by conjugation to the bs-40296G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Colon; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with MUM1 Monoclonal Antibody, Unconjugated (bsm-34186M) at 1:200 overnight at 4°C, followed by conjugation to the bs-40296G-HRP and DAB (C-0010) staining.