bs-10633R

[Primary Antibody]

BIOSS ANTIBODIES www.bioss.com.cn

CYT 19 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Applications: WB (1:500-2000)

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GenelD: 57412 SWISS: Q9HBK9

Target: CYT 19

Immunogen: KLH conjugated synthetic peptide derived from human CYT 19:

21-120/375.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Formation of methylated metabolites is a critical step in the

metabolism of inorganic arsenic. Arsenite methyltransferase (cyt19) is localized to the cytoplasm and operates in the transfer of a methyl group from AdoMet to trivalent arsenicals producing methylated and dimethylated arsenicals. It methylates arsenite to form methylarsonate which is reduced to methylarsonite. Methylarsonite acts as a substrate and is converted into a much less toxic compound dimethylarsinate. cyt19 is highly expressed in liver. Inherited variation in cyt19 may contribute to variation in

liver. Inherited variation in cyt19 may contribute to variation in arsenic metabolism and possibly arsenic-dependent

carcinogenesis in humans.

Reactivity: Mouse (predicted: Human,

Rat, Rabbit, Sheep, Cow, Chicken, Dog, Horse)


Predicted MW.: 42 kDa

Subcellular Nucleus Location:

VALIDATION IMAGES



Protein: heart(mouse) lysate at 40ug; Primary: rabbit Anti-CYT 19 (bs-10633R) at 1:300; Secondary: HRP conjugated Goat-Anti-rabbit lgG(bs-0295G-HRP) at 1: 5000; Predicted band size: 42 kD Observed band size: 42 kD



Sample: heart (Mouse) Lysate at 40 ug liver (Mouse) Lysate at 40 ug Primary: Anti-CYT 19 (bs-10633R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 42 kD Observed band size: 48 kD