


bsm-52813R**[Primary Antibody]****Thymidine kinase 1 Recombinant Rabbit mAb****BioSS**
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —**Host:** Rabbit**Isotype:** IgG**Clonality:** Recombinant**CloneNo.:** 9C1**GeneID:** 7083**SWISS:** P04183**Target:** Thymidine kinase 1**Immunogen:** A synthesized peptide derived from human Thymidine kinase: 2-38.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS(pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20 °C for one year. Avoid repeated freeze/thaw cycles.**Background:** Thymidine Kinase (TK1) is a highly conserved phosphotransferase that is present in most living cells. Thymidine Kinase catalyzes the phosphorylation reaction: deoxythymidine + ATP = deoxythymidine 5'-phosphate + ADP; it is thus involved in the reaction chain to introduce deoxythymidine into the DNA. Thymidine kinase is required for the action of many antiviral drugs, such as azidothymidine (AZT), and is also used to select hybridoma cell lines in the production of monoclonal antibodies. Thymidine Kinase has many clinical applications as it is only present in anticipation of cell division. Because of this, Thymidine Kinase can be used as a proliferation marker in the diagnosis, treatment, and follow-up of malignant diseases, especially hematological malignancies. Thymidine Kinase may be observed as a monomer, dimer, trimer or tetramer.**Applications:** WB (1:500-2000)**IHC-P** (1:50-100)**Flow-Cyt** (1:20-100)**ICC/IF** (1:50-100)**IP** (1:20-50)**Reactivity:** Human**Predicted**
MW.: 25 kDa**Subcellular**
Location: Cytoplasm**— VALIDATION IMAGES —**

Western blot analysis of Molt-4 cell lysate. Using Thymidine kinase 1 (bsm-52813R) monoclonal antibody at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.