

bs-6150R**[Primary Antibody]****GPR124 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

DATASHEET

| | | |
|---|---|--|
| Host: Rabbit Clonality: Polyclonal GeneID: 25960 Target: GPR124 Immunogen: KLH conjugated synthetic peptide derived from human GPR124: 1201-1303/1303. Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: Tumor endothelial markers (TEMs) are abundantly expressed in the blood vessels of human solid tumors during angiogenesis and neoangiogenesis. These include TEM1 (endosialin), TEM5 (G-protein coupled receptor 124) and TEM7 (plexin domain containing 1). TEMs are associated with the cell surface membrane at low levels in normal human and mouse tissues. TEM5 is a seven-pass transmembrane receptor, whereas TEM1, TEM7 and TEM8 span the membrane once. TEM5 expression is elevated during tumor angiogenesis and neoangiogenesis. TEM7 is highly expressed in tumor endothelium and neurons. Therefore, TEM5 and TEM7 may be suitable targets for the development of antiangiogenic therapies. | Isotype: IgG SWISS: Q96PE1 | Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) ICC/IF (1:100-500) ELISA (1:5000-10000) |
| | | Reactivity: Human, Mouse, Rat (predicted: Rabbit, Sheep, Cow, Dog, Horse) |
| | | Predicted MW.: 139 kDa |
| | | Subcellular Location: Cell membrane |

VALIDATION IMAGES

Sample: Lane 1: Placenta (Mouse) Lysate at 40 ug
Lane 2: Lung (Mouse) Lysate at 40 ug
Lane 3: Large intestine (Rat) Lysate at 40 ug
Lane 4: Lung (Rat) Lysate at 40 ug
Lane 5: U2os (Human) Cell Lysate at 30 ug
Primary: Anti-GPR124 (bs-6150R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 180-200 kD
Observed band size: 245 kD