

bs-2972R**[Primary Antibody]****HES1 Rabbit pAb****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

— DATASHEET —

Host: Rabbit	Isotype: IgG	Applications: IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Flow-Cyt (1 μ g/Test) ICC/IF (1:50-200) Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Cow, Horse) Predicted MW.: 30 kDa Subcellular Location: Nucleus
Clonality: Polyclonal		
GeneID: 3280	SWISS: Q14469	
Target: HES1		
Immunogen: KLH conjugated synthetic peptide derived from human HES1: 41-150/280.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Transcriptional repressor of genes that require a bHLH protein for their transcription. May act as a negative regulator of myogenesis by inhibiting the functions of MYOD1 and ASH1. Binds DNA on N-box motifs: 5'-CACNAG-3' with high affinity and on E-box motifs: 5'-CANNTG-3' with low affinity.		

— VALIDATION IMAGES —


Paraformaldehyde-fixed, paraffin embedded (human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HES1) Polyclonal Antibody, Unconjugated (bs-2972R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HES1) Polyclonal Antibody, Unconjugated (bs-2972R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



MCF-7 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (HES1) polyclonal Antibody, Unconjugated (bs-2972R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



MCF-7 cell; 4% Paraformaldehyde-fixed; Triton X-100 at room temperature for 20 min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Antibody incubation with (HES1) polyclonal Antibody, Unconjugated (bs-2972R) 1:100, 90 minutes at 37°C; followed by a conjugated Goat Anti-Rabbit IgG antibody at 37°C for 90 minutes, DAPI (blue, C02-04002) was used to stain the cell nuclei.



Blank control: Molt4. Primary Antibody (green line): Rabbit Anti-HES1 antibody (bs-2972R) Dilution: 1 μ g / 10⁶ cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody : Goat anti-rabbit IgG-AF647 Dilution: 1 μ g /test. Protocol The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

5%BSA to block non-specific protein-protein interactions for 30 min at room temperature
.Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature.
Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=5.7]** Hangbin Xia. et al. Combined magnesium and silicon ions synergistically promote functional regeneration of skeletal muscle by regulating satellite cell fate..Regenerative Biomaterials.2025 Feb 19:12:rbaf008. IF ;Mouse. 40130074
- **[IF=5.17]** Wang et al. Hes1 triggers epithelial-mesenchymal transition (EMT)-like cellular marker alterations and promotes invasion and metastasis of nasopharyngeal carcinoma by activating the PTEN/AKT pathway. (2015) Oncotarge. 6:36713-30 WB,IHC ;Human. 26452025
- **[IF=5.008]** Gao et al. Hes1 promotes cell proliferation and migration by activating Bmi-1 and PTEN/Akt/GSK3 β pathway in human colon cancer. (2015) Oncotarge. 6:38667-80 IHC,ICC,WB ;Human, Mouse. 26452029
- **[IF=3.6]** Yahao Zhao. et al. PEAR1 Promotes Myoblast Proliferation Through Notch Signaling Pathway. BIOLOGY-BASEL. 2024 Dec;13(12):1063 WB ;Mouse. 39765730
- **[IF=4.486]** Chong Yin. et al. MACF1 alleviates aging-related osteoporosis via HES1. 2021 Jun 15 WB,IHC ;Mouse. 34133068