

bs-6177R

[Primary Antibody]

RAB9 Rabbit pAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


DATASHEET

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500) Reactivity: Human, Rat (predicted: Mouse, Pig, Sheep, Cow, Chicken, Dog) Predicted MW.: 23 kDa Subcellular Location: Cell membrane ,Cytoplasm
Clonality: Polyclonal		
GeneID: 9367	SWISS: P51151	
Target: RAB9		
Immunogen: KLH conjugated synthetic peptide derived from human RAB9: 65-170/201.		
Purification: affinity purified by Protein A		
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: RAB proteins are GTPases that regulate vesicular trafficking and reside in specific intracellular compartments. RAB9 has been localized to components of the endocytic/exocytic pathway. It has been implicated in the recycling of membrane receptors, such as the mannose 6-phosphate receptor from early endosomes to the trans Golgi network.		


VALIDATION IMAGES




Sample: K562(Human) Cell Lysate at 30 ug
Primary: Anti-RAB9 (bs-6177R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at
1/20000 dilution Predicted band size: 23 kD
Observed band size: 23 kD



Sample: HeLa(Human) Cell Lysate at 30 ug
Primary: Anti-RAB9 (bs-6177R) at 1/500 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at
1/20000 dilution Predicted band size: 23 kD
Observed band size: 23 kD



Tissue/cell: rat brain tissue; 4%
Paraformaldehyde-fixed and paraffin-
embedded; Antigen retrieval: citrate buffer (
0.01M, pH 6.0), Boiling bathing for 15min; Block
endogenous peroxidase by 3% Hydrogen
peroxide for 30min; Blocking buffer (normal goat
serum,C-0005) at 37°C for 20 min; Incubation:
Anti-RAB9 Polyclonal Antibody,
Unconjugated(bs-6177R) 1:200, overnight at 4°C,
followed by conjugation to the secondary
antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded
(Rat stomach); Antigen retrieval by boiling in
sodium citrate buffer (pH6.0) for 15min; Block
endogenous peroxidase by 3% hydrogen
peroxide for 20 minutes; Blocking buffer (normal
goat serum) at 37°C for 30min; Antibody
incubation with (RAB9) Polyclonal Antibody,
Unconjugated (bs-6177R) at 1:400 overnight at

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

4°C, followed by operating according to SP
Kit(Rabbit) (sp-0023) instructions and DAB
staining.