## bs-1066R

## [ Primary Antibody ]

# NME1/NM23A Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

GeneID: 4830 **SWISS:** P15531

Target: NME1/NM23A

Immunogen: KLH conjugated synthetic peptide derived from human Nm23-H1:

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: This gene (NME1) was identified because of its reduced mRNA

transcript levels in highly metastatic cells. Nucleoside diphosphate kinase (NDK) exists as a hexamer composed of 'A' (encoded by this gene) and 'B' (encoded by NME2) isoforms. Mutations in this gene have been identified in aggressive neuroblastomas. Two transcript variants encoding different isoforms have been found for this gene. Co-transcription of this gene and the neighboring downstream gene (NME2) generates naturally-occurring transcripts (NME1-NME2), which encodes a fusion protein comprised of sequence

sharing identity with each individual gene product.

Applications: WB (1:500-2000)

**IHC-P** (1:100-500) **IHC-F** (1:100-500) **IF** (1:100-500) Flow-Cyt (1µg/Test)


Reactivity: Human, Mouse, Rat

(predicted: Rabbit, Pig, Cow, Dog, Horse)


Predicted MW.: 17 kDa

Subcellular Cytoplasm ,Nucleus


### - VALIDATION IMAGES -




Sample: Hela(Human) Cell Lysate at 30 ug Primary: Anti-NM23A (bs-1066R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 17 kD Observed band size: 18 kD



Tissue/cell: mouse heart tissue: 4% Paraformaldehyde-fixed and paraffinembedded: Antigen retrieval: citrate buffer ( 0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-NME1/Nm23-H1/NDKA Polyclonal Antibody, Unconjugated(bs-1066R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Blank control: RSC96(blue). Primary Antibody:Rabbit Anti-NME1 antibody(bs-1066R), Dilution: 1µg in 100 µL 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions ); Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA. Protocol The cells were fixed with 2% paraformaldehyde (10 min), then permeabilized with 90% ice-cold methanol for 30 min on ice. Antibody (bs-1066R, 1μg /1x10^6 cells) were incubated for 30 min on the ice, followed by 1 X PBS containing 0.5% BSA + 10% goat serum (15 min) to block non-specific protein-protein interactions. Then the Goat Antirabbit IgG/PE antibody was added into the blocking buffer mentioned above to react with the primary antibody of bs-1066R at 1/200 dilution for 30 min on ice. Acquisition of 20,000 events was performed.



Blank control (blue line): A549 (blue). Primary Antibody (green line): Rabbit Anti-NME1 antibody (bs-1066R) Dilution: 1µg/10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG . Secondary Antibody (white blue line): Goat anti-rabbit IgG-PE Dilution: 1μg /test. Protocol The cells were fixed with 2% paraformaldehyde (10 min) , then permeabilized with 90% ice-cold methanol for 30 min on ice. Cells stained with Primary Antibody for 30 min at room temperature. The cells were then incubated in  $\ensuremath{\mathbf{1}}$ X PBS/2%BSA/10% goat serum to block nonspecific protein-protein interactions followed by the antibody for 15 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.