

bsm-63079R**[Primary Antibody]**

BioSS
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Phospho-mTOR(S2481) Recombinant Rabbit mAb**— DATASHEET —**


Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)
Clonality: Recombinant	CloneNo.: 15D14	
GeneID: 2475	SWISS: P42345	
Target: Phospho-mTOR(S2481)		
Immunogen: A synthesized peptide derived from human mTOR around the phosphorylation site of S2481: IH-pS-FI(全长2549aa).		
Purification: affinity purified by Protein A		Reactivity: Human, Mouse, Rat Predicted MW.: 289 kDa Subcellular Location: Cell membrane ,Cytoplasm
Concentration: 1mg/ml		
Storage: 10mM phosphate buffered saline , pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: mTOR is one of a family of proteins involved in cell cycle progression, DNA recombination, and DNA damage detection. In rat, it is a 289-kDa protein (symbolized RAFT1) with significant homology to the Saccharomyces cerevisiae protein TOR1 and has been shown to associate with the immunophilin FKBP12 in a rapamycin dependent fashion. The FKBP12-rapamycin complex is known to inhibit progression through the G1 cell cycle stage by interfering with mitogenic signaling pathways involved in G1 progression in several cell types, as well as in yeast. The binding of FRAP to FKBP12-rapamycin correlated with the ability of these ligands to inhibit cell cycle progression.		

— VALIDATION IMAGES —

293T (H) cells were treated with or without Calyculin A (100 nM) for 60 min, 25 µg total protein per lane of cell lysates (see on figure) probed with Phospho-mTOR(S2481) monoclonal antibody, unconjugated (bsm-63079R) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Breast Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Phospho-mTOR(S2481) Monoclonal Antibody, Unconjugated(bsm-63079R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.



Paraformaldehyde-fixed, paraffin embedded Human Lung Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Phospho-mTOR(S2481) Monoclonal Antibody, Unconjugated(bsm-63079R) at 1:200 overnight at 4°C, followed by conjugation to the bs-0295G-HRP and DAB (C-0010) staining.