bs-17508R

[Primary Antibody]

Bioss ANTIBODIES

SKOR1 Rabbit pAb

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

IHC-F (1:100-500)

(predicted: Human, Rabbit, Pig, Sheep, Cow, Horse)

IF (1:100-500)

100 kDa

Applications: IHC-P (1:100-500)

Reactivity: Mouse, Rat

Predicted

MW.:

Subcellular Nucleus

- DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal GeneID: 390598 Target: SKOR1

Immunogen: KLH conjugated synthetic peptide derived from human SKOR1:

1-100/965.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated


freeze/thaw cycles.

mutations to chromosome 15.


Background: LBXCOR1 is a 965 amino acid protein belonging to the SKI family.

Localizing to nucleus, LBXCOR1 is highly expressed in the central nervous system (CNS) as well as developing spinal cord and adult brain and testis. LBXCOR1 contains a CH1 domain which is required for transcriptional repression and also acts as a transcriptional co-repressor of LBX1. LBXCOR1 additionally interacts with SMAD1, SMAD2 and SMAD3, and inhibits BMP signaling. Existing as three alternatively spliced isoforms, the gene encoding LBXCOR1 maps to human chromosome 15q23. Encoding more than 700 genes, chromosome 15 is made up of approximately 106 million base pairs and consists of about 3% of the human genome. Angelman and Prader-Willi syndromes, as well as Tay-Sachs disease and Marfan syndorme, are all associated with

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SKOR1) Polyclonal Antibody, Unconjugated (bs-17508R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SKOR1) Polyclonal Antibody, Unconjugated (bs-17508R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.