

bs-41334R

[Primary Antibody]

ASC/TMS1 Rabbit pAb

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800

DATASHEET

Host: Rabbit	Isotype: IgG	Applications: WB (1:500-2000)
Clonality: Polyclonal		Reactivity: Mouse, Rat
Target: ASC/TMS1		
Immunogen: Recombinant mouse PYCARD protein: 1-193/193.		Predicted MW.: 22 kDa
Purification: affinity purified by Protein A		Subcellular Location: Cytoplasm ,Nucleus
Concentration: 1mg/ml		
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.		
Background: Promotes caspase-mediated apoptosis. This proapoptotic activity is mediated predominantly through the activation of caspase-9. May be a component of the inflammasome, a protein complex which also includes NALP2, CARD8 and CASP1 and whose function would be the activation of proinflammatory caspases. Tissue specificity: Widely expressed at low levels. Detected in peripheral blood leukocytes, lung, small intestine, spleen, thymus, colon and at lower levels in placenta, liver and kidney. Very low expression in skeletal muscle, heart and brain. Detected in the leukemia cell lines HL-60 and U937, but not in Jurkat T-cell lymphoma and Daudi Burkitt's lymphoma.		

VALIDATION IMAGES



Sample: Lane 1: Mouse Lymph node tissue lysates
Lane 2: Mouse Lung tissue lysates
Lane 3: Mouse Spleen tissue lysates
Lane 4: Rat Lymph node tissue lysates
Lane 5: Rat Lung tissue lysates
Lane 6: Rat Spleen tissue lysates
Primary: Anti-ASC/TMS1 (bs-41334R) at 1/1000 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 22 kDa
Observed band size: 24 kDa

SELECTED CITATIONS

- [IF=8.8]** Egidi Maren Juliane. et al. Anti-angiogenic therapy using the multi-tyrosine kinase inhibitor Regorafenib enhances tumor progression in a transgenic mouse model of β -cell carcinogenesis. BRIT J CANCER. 2023 Aug;;1-13 IHC ;Mouse. 37620408
- [IF=7]** Yueqi Yang. et al. DHA and EPA Alleviate Epileptic Depression in PTZ-Treated Young Mice Model by Inhibiting Neuroinflammation through Regulating Microglial M2 Polarization and Improving Mitochondrial Metabolism. ANTIOXIDANTS-BASEL. 2023 Dec;12(12):2079 WB ;Mouse. 38136199