## bs-1317R

# [ Primary Antibody ]

# Bioss

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

**IHC-P** (1:100-500)

**IHC-F** (1:100-500)

ICC/IF (1:100-500)

Mouse, Pig, Sheep, Cow,

**IF** (1:100-500) **Flow-Cyt** (1ug/Test)

Reactivity: Rat (predicted: Human,

Dog, Horse)

Subcellular Location: Cytoplasm ,Nucleus

Applications: WB (1:500-2000)

Predicted MW.: 82 kDa

# STAT1 Rabbit pAb

- DATASHEET -

**Host:** Rabbit **Isotype:** IgG

Clonality: Polyclonal

**GenelD:** 6772 **SWISS:** P42224

Target: STAT1

**Immunogen:** KLH conjugated synthetic peptide derived from human STAT1:

251-380/750.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated


freeze/thaw cycles.

**Background:** The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and


31C. [provided by RefSeq, Jun 2020]

growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. The protein encoded by this gene can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens. The protein plays an important role in immune responses to viral, fungal and mycobacterial pathogens. Mutations in this gene are associated with Immunodeficiency 31B, 31A, and

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (Rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (STAT1) Polyclonal Antibody, Unconjugated (bs-1317R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Tissue/cell: rat brain tissue; 4%
Paraformaldehyde-fixed and paraffinembedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min; Incubation: Anti-STAT1 Polyclonal Antibody, Unconjugated(bs-1317R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

### - SELECTED CITATIONS -

- [IF=17.4] Zhesheng He. et al. The therapeutic effect of gold clusters in an EAE model of multiple sclerosis by regulating T cell differentiation. NANO TODAY. 2024 Feb;54:102128 WB; Mouse. 10.1016/j.nantod.2023.102128
- [IF=7.9] Tiantian Wang. et al. Baicalein alleviates cardiomyocyte death in EAM mice by inhibiting the JAK-STAT1/4 signalling pathway. PHYTOMEDICINE. 2024 Jun;128:155558 WB; Mouse. 38547614
- [IF=5.58] Zhou, Zhiwei, et al. "microRNA let-7c is essential for the anisomycin-elicited apoptosis in Jurkat T cells by linking JNK1/2 to AP-1/STAT1/STAT3 signaling." Scientific Reports 6 (2016): 24434. WB; Human. 27087117
- [IF=5.714] Chenglin Li. et al. Ranitidine as an adjuvant regulates macrophage polarization and activates CTLs through the PI3K-Akt2 signaling pathway. INT IMMUNOPHARMACOL. 2023 Mar;116:109729 WB; Mouse. 10.1016/j.intimp.2023.109729
- [IF=6.304] Chen Y et al. Dendritic cells-derived interferon-λ1 ameliorated inflammatory bone destruction through inhibiting osteoclastogenesis. Cell Death Dis. 2020 Jun 2;11(6):414. WB; Mouse. 32488049