bs-9308R

- DATASHEET -

[Primary Antibody]

KATNA1 Rabbit pAb



www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Applications: WB (1:500-2000) IHC-P (1:100-500) IHC-F (1:100-500) IF (1:50-200)

Reactivity: Human, Mouse, Rat (predicted: Rabbit, Pig, Sheep, Cow, Chicken, Dog, Horse)

Predicted MW.: ^{56 kDa}

Subcellular Location: Cytoplasm

Host: Rabbit Clonality: Polyclonal GenelD: 11104

SWISS: 075449

Isotype: IgG

Target: KATNA1

Immunogen: KLH conjugated synthetic peptide derived from human KATNA1/Katanin p60 A1: 221-320/491.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Microtubules are polymers of alpha and beta subunits that form the mitotic spindle and assist in the organization of membranous organelles during interphase. Katanin p60 A1, also known as KATNA1, is a 491 amino acid protein that belongs to the AAA ATPase family and is involved in microtubule regulation. Localized to the cytoplasm and to the centrosome, Katanin p60 A1 functions to sever and disassemble microtubules in an ATP-dependent manner, thus promoting the rapid reorganization of cellular microtubule arrays and playing an important role in microtubule release from the centrosome after nucleation. Katanin p60 A1, which exists as two alternatively spliced isoforms, can homooligomerize into hexameric rings whose activity is stimulated by the presence of microtubules.

- VALIDATION IMAGES



Sample: Fetal brain (Mouse) Lysate at 40 ug Primary: Anti-KATNA1 (bs-9308R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size:56 kD Observed band size: 56 kD



Sample: Lane 1: Mouse Cerebrum tissue lysates Lane 2: Mouse Liver tissue lysates Lane 3: Rat Cerebrum tissue lysates Lane 4: Rat Liver tissue lysates Lane 5: Human HeLa cell lysates Lane 6: Human MCF-7 cell lysates Lane 7: Human Jurkat cell lysates Primary: Anti-KATNA1 (bs-9308R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 56 kDa Observed band size: 50 kDa



Paraformaldehyde-fixed, paraffin embedded (rat stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (KATNA1) Polyclonal Antibody, Unconjugated (bs-9308R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KATNA1) Polyclonal Antibody, Unconjugated (bs-9308R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.