bs-9081R

[Primary Antibody]

www.bioss.com.cn

sales@bioss.com.cn techsupport@bioss.com.cn

DATASHEET -

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

AKIRIN1 Rabbit pAb

GenelD: 79647 SWISS: Q9H9L7

Target: AKIRIN1

Immunogen: KLH conjugated synthetic peptide derived from human AKIRIN1:

101-192/192.

Purification: affinity purified by Protein A

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: AKIRIN1 is dispensable in the mouse, and neither knockout mice nor cells derived from them have obvious distinctive phenotypes. In contrast, Akirin2 is required for development in the mouse and knockout of both Akirin homologs in mice show that Akirin2 is required downstream of toll-like receptor (TLR), TNF-alpha and IL-1beta signaling, and for the production of IL-6. Akirin2 is functionally closer to the single gene in Drosophila, as the homozygous null D. melanogaster Akirin mutants show a similar, mid-to-early embryonic death. The highly conserved, nuclearlocalized AKIRIN1 and Akirin2 proteins critically regulate the transcription of NF-kB dependent genes and are required for defense against Gram-negative bacteria in the immune deficiency and NF-kB pathways.

Applications: IHC-P (1:100-500)

400-901-9800

IHC-F (1:100-500) **IF** (1:100-500)


Reactivity: Rat (predicted: Human,

Mouse, Sheep, Dog, Horse)


Predicted MW.:

Subcellular Nucleus

VALIDATION IMAGES



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (AKIRIN1) Polyclonal Antibody, Unconjugated (bs-9081R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (AKIRIN1) Polyclonal Antibody, Unconjugated (bs-9081R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.