


HSD17B1 Rabbit pAb

Catalog Number: bs-6603R
Target Protein: HSD17B1
Concentration: 1mg/ml
Form: Liquid
Host: Rabbit
Clonality: Polyclonal
Isotype: IgG
Applications: WB (1:500-2000)
Reactivity: Mouse (predicted:Human, Rat, Rabbit, Pig, Cow, Dog, Horse)
Predicted MW: 36 kDa
Entrez Gene: 3292
Swiss Prot: P14061
Source: KLH conjugated synthetic peptide derived from human HSD17B1: 21-120/328.
Purification: affinity purified by Protein A
Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Background: This gene encodes a member of the 17beta-hydroxysteroid dehydrogenase family of short-chain dehydrogenases/reductases. It has a dual function in estrogen activation and androgen inactivation and plays a major role in establishing the estrogen E2 concentration gradient between serum and peripheral tissues. The encoded protein catalyzes the last step in estrogen activation, using NADPH to convert estrogens E1 and E2 and androgens like 4-androstenedione, to testosterone. It has an N-terminal short-chain dehydrogenase domain with a cofactor binding site, and a narrow, hydrophobic C-terminal domain with a steroid substrate binding site. This gene is expressed primarily in the placenta and ovarian granulosa cells, and to a lesser extent, in the endometrium, adipose tissue, and prostate. Polymorphisms in this gene have been linked to breast and prostate cancer. A pseudogene of this gene has been identified. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2016]

VALIDATION IMAGES



Sample: Placenta (Mouse) Lysate at 40 ug Primary: Anti-HSD17B1 (bs-6603R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 36 kD Observed band size: 38 kD



Sample: Ovary (Mouse) Lysate at 40 ug Primary: Anti-HSD17B1 (bs-6603R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 36 kD Observed band size: 36 kD

PRODUCT SPECIFIC PUBLICATIONS

[IF=8.469] Xiaofeng Zhou. et al. DNA methylation mediated RSPO2 to promote follicular development in mammals. Cell Death Dis. 2021 Jun;12(7):1-13 WB ; Pig . 34175894