


bsm-60349R**[Primary Antibody]****Bioss**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn


400-901-9800

Pan Actin Recombinant Rabbit mAb**DATASHEET****Host:** Rabbit**Clonality:** Recombinant**Target:** Pan Actin**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.**Background:** Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells. In vertebrates 3 main groups of actin isoforms, alpha, beta and gamma have been identified. The alpha actins are found in muscle tissues and are a major constituent of the contractile apparatus. The beta and gamma actins coexist in most cell types as components of the cytoskeleton and as mediators of internal cell motility. Actin proteins have very high homology. The immunogen for this product is based on the ACTC1 protein but this antibody will also recognize: ACTB, ACTA1, ACTG1, ACTA2, ACTG2.**Isotype:** IgG**CloneNo.:** F3C6**Applications:** WB (1:500-1:1000)**IHC-P** (1:100-500)**IHC-F****IF****Flow-Cyt** (1:50-1:100)**ICC/IF** (1:50)**Reactivity:** Human, Mouse, Rat**Predicted MW.:** 42 kDa**Subcellular Location:** Cell membrane**VALIDATION IMAGES**


Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:1000 Primary ab incubation condition: 2 hours at room temperature
Secondary ab: Goat Anti-Rabbit IgG H&L (HRP)
Lysate: HeLa, Jurkat, HEK293, NIH/3T3, C2C12, Mouse brain, Rat brain Protein loading quantity: 20 µg Exposure time: 60 s Predicted MW: 42 kDa
Observed MW: 42 kDa




Paraformaldehyde-fixed, paraffin embedded (mouse stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (Pan Actin) Monoclonal Antibody, Unconjugated (bsm-60349R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (human heart); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation with (Pan Actin) Monoclonal Antibody, Unconjugated (bsm-60349R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Cell line: NIH/3T3 Fixative: 4% Paraformaldehyde Permeabilization: 0.1% TritonX-100 Primary ab dilution: 1:50 Primary incubation condition: 4°C overnight Secondary ab: Goat Anti-Rabbit IgG Nuclear counter stain: DAPI (Blue) Comment: Color green is the positive signal for bsm-60349R



Cell line: A431 Fixative: 100% Ice-cold methanol Permeabilization: 0.1% TritonX-100 Primary ab dilution: 1:50 Primary incubation condition: 4°C overnight Secondary ab: Goat Anti-Rabbit IgG Nuclear counter stain: DAPI (Blue) Comment: Color green is the positive signal for bsm-60349R



Cell line: Jurkat Fixative: 4% Paraformaldehyde Permeabilization: 90% Methanol Primary ab dilution: 1:100 Secondary ab: Goat anti Rabbit IgG Unlabelled control: The cell without incubation with primary antibody and secondary antibody (Black line). Isotype control: Rabbit monoclonal IgG (Blue line). Comment: Line red is the positive signal for bsm-60349R

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.