bsm-60091M

[Primary Antibody]

BIOSS

www.bioss.com.cn sales@bioss.com.cn techsupport@bioss.com.cn 400-901-9800

Di-Methyl-Histone H3 (Lys36) Mouse mAb

DATASHEET -

Host: Mouse Isotype: IgG1
Clonality: Monoclonal CloneNo.: B6B6

Target: Di-Methyl-Histone H3 (Lys36) **Purification:** affinity purified by Protein G

Concentration: 1mg/ml

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50%

Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated

freeze/thaw cycles.

Background: Modulation of the chromatin structure plays an important role in

the regulation of transcription in eukaryotes. The nucleosome, made up of four core histone proteins (H2A, H2B, H3 and H4), is the primary building block of chromatin. The N-terminal tail of core histones undergoes different posttranslational modifications including acetylation, phosphorylation and methylation. These modifications occur in response to cell signal stimuli and have a direct effect on gene expression. In most species, the histone H2B is primarily acetylated at lysines 5, 12, 15 and 20. Histone H3 is primarily acetylated at lysines 9, 14, 18 and 23. Acetylation at lysine 9 appears to have a dominant role in histone deposition and chromatin assembly in some organisms. Phosphorylation at Ser10 of histone H3 is tightly correlated with chromosome condensation during both mitosis and meiosis.


Applications: WB (1:500-1000)

IHC-P (1:100-500) IHC-F (1:100-500) IF (1:100-500)


Reactivity: Human, Mouse, Rat

Subcellular Nucleus


VALIDATION IMAGES



Blocking buffer: 5% NFDM/TBST Primary ab dilution: 1:1000 Primary ab incubation condition: 4°C overnight Secondary ab: Goat Anti-Mouse IgG H&L (HRP) Lysate: HeLa, Neuro-2a, BRL, Mouse spleen, Mouse kidney Protein loading quantity: 20 µg Exposure time: 30 s Predicted MW: 15 kDa Observed MW: 15 kDa



Tissue: Human colon cancer Section type:
Formalin fixed & Paraffin -embedded section
Retrieval method: High temperature and high
pressure Retrieval buffer: Tris/EDTA buffer, pH
9.0 Primary ab dilution: 1:100 Primary ab
incubation condition: 1 hour at room
temperature Secondary ab: SP
Kit(Mouse)(sp-0024) Counter stain: Hematoxylin
(Blue) Comment: Color brown is the positive
signal for bsm-60091M



Tissue: Rat testis Section type: Formalin fixed & Paraffin -embedded section Retrieval method: High temperature and high pressure Retrieval buffer: Tris/EDTA buffer, pH 9.0 Primary ab dilution: 1:100 Primary ab incubation condition: 1 hour at room temperature Secondary ab: SP Kit(Mouse)(sp-0024) Counter stain: Hematoxylin (Blue) Comment: Color brown is the positive signal for bsm-60091M