

bs-3397R**[Primary Antibody]****phospho-SGK1 (Thr256) Rabbit pAb**

Bioss
ANTIBODIES

www.bioss.com.cn

sales@bioss.com.cn


techsupport@bioss.com.cn

400-901-9800


DATASHEET**Host:** Rabbit**Isotype:** IgG**Clonality:** Polyclonal**GeneID:** 6446**SWISS:** O00141**Target:** SGK1 (Thr256)**Immunogen:** KLH conjugated Synthesised phosphopeptide derived from human SGK1 around the phosphorylation site of Thr256: TS(p-T)FC.**Purification:** affinity purified by Protein A**Concentration:** 1mg/ml**Storage:** 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.


Background: SGK1 is a protein kinase that plays an important role in cellular stress response. SGK1 activates certain potassium, sodium, and chloride channels, suggesting an involvement in the regulation of processes such as cell survival, neuronal excitability, and renal sodium excretion. Sustained high levels of SGK1 and activity may contribute to conditions such as hypertension and diabetic nephropathy. This protein also mediates cell survival signals, as it has been shown to phosphorylate and negatively regulate the pro apoptotic FOXO3A protein. Ser 422 is a critical site on the protein and may be involved in its activation.

Applications: WB (1:500-2000)**IHC-P** (1:100-500)**IHC-F** (1:100-500)**IF** (1:100-500)**Flow-Cyt** (2ug/Test)**ICC/IF** (1:25)**Reactivity:** Human, Mouse
(predicted: Rat)**Predicted
MW.:** 49 kDa**Subcellular
Location:** Cell membrane ,Cytoplasm
Nucleus**VALIDATION IMAGES**


Sample: Pancreas (Mouse) Lysate at 40 ug
Primary: Anti-Phospho-SGK1 (Thr256) (bs-3397R)
at 1/1000 dilution Secondary: IRDye800CW Goat
Anti-Rabbit IgG at 1/20000 dilution Predicted
band size: 49 kD Observed band size: 51 kD



Ti Tissue/cell: human lung cancer; 4%
Paraformaldehyde-fixed and paraffin-
embedded; Antigen retrieval: citrate buffer (
0.01M, pH 6.0), Boiling bathing for 15min; Block
endogenous peroxidase by 3% Hydrogen
peroxide for 30min; Blocking buffer (normal goat
serum, C-0005) at 37°C for 20 min; Incubation:
Anti-Phospho-SGK1 (Thr256) Polyclonal
Antibody, Unconjugated (bs-3397R) 1:200,
overnight at 4°C, followed by conjugation to the
secondary antibody (SP-0023) and DAB (C-0010)
staining



Hela cell; 4% Paraformaldehyde-fixed; Triton
X-100 at room temperature for 20 min; Blocking
buffer (normal goat serum, C-0005) at 37°C for 20
min; Antibody incubation with (Phospho-SGK1
(Thr256)) polyclonal Antibody, Unconjugated
(bs-3397R) 1:25, 90 minutes at 37°C; followed by
a conjugated Goat Anti-Rabbit IgG antibody at
37°C for 90 minutes, DAPI (blue, C02-04002) was
used to stain the cell nuclei.



Blank control: Mouse spleen. Primary Antibody
(green line): Rabbit Anti-SGK1 antibody
(bs-3397R) Dilution: 2µg / 10⁶ cells; Isotype

Important Note: This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Control Antibody (orange line): Rabbit IgG .
Secondary Antibody : Goat anti-rabbit IgG-PE
Dilution: 1µg /test. Protocol The cells were fixed with 4% PFA (10min at room temperature)and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.

— SELECTED CITATIONS —

- **[IF=3.288]** Chai D et al. β 2-microglobulin has a different regulatory molecular mechanism between ER+ and ER- breast cancer with HER2.BMC Cancer. 2019 Mar 12;19(1):223. WB,IHC ;Human. 30866857