
FAK Rabbit pAb

Catalog Number: bs-1340R

Target Protein: FAK

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000), IHC-P (1:100-500), IHC-F (1:100-500), IF (1:100-500)

Reactivity: Human, Mouse, Rat

Predicted MW: 116 kDa

Subcellular Cell membrane ,Cytoplasm ,Nucleus

Locations:

Entrez Gene: 5747

Swiss Prot: Q05397

Source: KLH conjugated synthetic peptide derived from human FAK: 901-1052/1052.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.


Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: Non-receptor protein-tyrosine kinase implicated in signaling pathways involved in cell motility, proliferation and apoptosis. Activated by tyrosine-phosphorylation in response to either integrin clustering induced by cell adhesion or antibody cross-linking, or via G-protein coupled receptor (GPCR) occupancy by ligands such as bombesin or lysophosphatidic acid, or via LDL receptor occupancy. Plays a potential role in oncogenic transformations resulting in increased kinase activity. [SUBCELLULAR LOCATION] Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Note=Constituent of focal adhesions.


VALIDATION IMAGES




Sample: Lane 1: Cerebrum (Mouse) Lysate at 40 ug Lane 2: NIH/3T3 (Mouse) Cell Lysate at 30 ug Lane 3: Cerebrum (Rat) Lysate at 40 ug Lane 4: 293T (Human) Cell Lysate at 30 ug Lane 5: A431 (Human) Cell Lysate at 30 ug Lane 6: Huvec (Human) Cell Lysate at 30 ug Lane 7: A549 (Human) Cell Lysate at 30 ug Primary: Anti-FAK (bs-1340R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 125 kD Observed band size: 120 kD




Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAK) Polyclonal Antibody, Unconjugated (bs-1340R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAK) Polyclonal Antibody, Unconjugated (bs-1340R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAK) Polyclonal Antibody, Unconjugated (bs-1340R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: mouse lung tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min; Incubation: Anti-FAK/PTK2 Polyclonal Antibody, Unconjugated (bs-1340R) 1:200, overnight at 4°C; The secondary antibody was Goat Anti-Rabbit IgG, Cy3 conjugated (bs-0295G-Cy3) used at 1:200 dilution for 40 minutes at 37°C. DAPI (5ug/ml, blue, C-0033) was used to stain the cell nuclei

PRODUCT SPECIFIC PUBLICATIONS

[IF=11.161] Huang, Chengmei. et al. Tumor cell-derived SPON2 promotes M2-polarized tumor-associated macrophage infiltration and cancer progression by activating PYK2 in CRC. J Exp Clin Canc Res. 2021 Dec;40(1):1-17 WB ; human . 34583750

[IF=7.59] Zhenyin Chen. et al. Dynamic stiffening collagen-coated substrate enhances osteogenic differentiation of mesenchymal stem cells through integrin $\alpha 2 \beta 1$. BIOMATER SCI-UK. 2023 May; WB, IF, ICC ; Rat . 37233200

[IF=4.9] Mohammad Abul Hasnat. et al. Action Mechanisms of Exosomes Derived from GD3/GD2-Positive Glioma Cells in the Regulation of Phenotypes and Intracellular Signaling: Roles of Integrins. INT J MOL SCI. 2024 Jan;25(23):12752 WB ; Human . 39684463

[IF=4.848] Danning Wang. et al. Cyclin G2 Inhibits Oral Squamous Cell Carcinoma Growth and Metastasis by Binding to IGFBP3 and Regulating the FAK-SRC-STAT Signaling Pathway. Front Oncol. 2020; 10: 560572 WB, IP ; Human . 33240810

[IF=5.307] Jian Song. et al. The dual FAK-HDAC inhibitor MY-1259 displays potent activities in gastric cancers in vitro and in vivo. BIOORG CHEM. 2023 Feb;131:106328 WB ; Human . 36542986