

bs-41214R**[Primary Antibody]****NCR1 Rabbit pAb****BioSS**
ANTIBODIES

www.bioss.com.cn


sales@bioss.com.cn

techsupport@bioss.com.cn

400-901-9800

— DATASHEET —

| | | |
|--|---|--|
| Host: Rabbit Clonality: Polyclonal GeneID: 9437 Target: NCR1 Purification: affinity purified by Protein A Concentration: 1mg/ml Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. Background: The natural cytotoxicity receptors (NCRs) are a recently characterized family of Ig-like activation receptors that appear to be major triggering receptors in tumor cell recognition. NCR1 is a glycoprotein that has two extracellular Ig-like domains followed by a ~40 amino acid residue stalk region, a type I transmembrane domain, and a short cytoplasmic tail. NCR1 has been shown to represent a novel NK cell-specific molecule involved in human NK cell activation. NCR1 has been implicated in NK cell-mediated lysis of several autologous tumor cells and pathogen-infected cell lines. | Isotype: IgG SWISS: O76036 | Applications: WB (1:500-2000) Reactivity: Human Predicted MW.: 31 kDa Subcellular Location: Cell membrane |
|--|---|--|

— VALIDATION IMAGES —

Sample: Lane 1: Human HL60 cell lysates Lane 2:
 Human HL60 cell lysates Primary: Anti-NCR1
 (bs-41214R) at 1/1000 dilution Secondary:
 IRDye800CW Goat Anti-Rabbit IgG at 1/20000
 dilution Predicted band size: 31 kD Observed
 band size: 46 kD

— SELECTED CITATIONS —

- **[IF=15.304]** Yao Lei. et al. Phytochemical natural killer cells reprogram tumor microenvironment for potent immunotherapy of solid tumors. BIOMATERIALS. 2022 Jun;;121635 WB,IF,FCM ;Mouse.
 10.1016/j.biomaterials.2022.121635