

Phospho-LATS1 (Thr1079) Rabbit pAb

Catalog Number: bs-3245R

Target Protein: Phospho-LATS1 (Thr1079)

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:500-2000)

Reactivity: Human (predicted:Mouse, Rat)

Predicted MW: 124 kDa

Entrez Gene: 26524

Swiss Prot: Q9NRM7

Source: KLH conjugated synthesised phosphopeptide derived from human LATS1 around the phosphorylation site of Thr1079: EF(P-T)FR.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: This gene encodes a serine/threonine protein kinase belonging to the LATS tumor suppressor family. The protein localizes to centrosomes during interphase, and early and late metaphase. It interacts with the centrosomal proteins aurora-A and ajuba and is required for accumulation of gamma-tubulin and spindle formation at the onset of mitosis. It also interacts with a negative regulator of p53 and may function in a positive feedback loop with p53 that responds to cytoskeleton damage. Additionally, it can function as a co-repressor of androgen-responsive gene expression. [provided by RefSeq].

VALIDATION IMAGES



Sample: HeLa(Human) Cell Lysate at 30 ug Primary: Anti-Phospho-LATS1 (Thr1079) (bs-3245R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 124 kD
Observed band size: 124 kD

PRODUCT SPECIFIC PUBLICATIONS

[IF=5.5] Li Lin. et al. GPR137 inactivates Hippo signaling to promote gastric cancer cell malignancy. BIOL DIRECT. 2024 Dec;19(1):1-16
WB,CoIP ; Human . 38163861

[IF=4.5] Shouying Xu. et al. ARID1A restrains EMT and stemness of ovarian cancer cells through the Hippo pathway. INT J ONCOL. 2024
Aug;65(2):1-11 WB ; Human . 38873993

[IF=4.7] Tianwei Dong. et al.Cannabidiol Ameliorates Doxorubicin-Induced Myocardial Injury via Activating Hippo Pathway.DRUG DESIGN
DEVELOPMENT AND THERAPY.2025 Jan 24;19:569-583. Western blot ; Rat, mouse . 39876987

[IF=1.4] Bingyi Chen. et al. The inhibition of γ -Aminobutyric Acid B1 receptor regulates angiogenesis via the Hippo/YAP signaling pathway.
ANN VASC SURG. 2024 Jul; WB ; Human . 39025214