
FGFR3 Rabbit pAb

Catalog Number: bs-1301R

Target Protein: FGFR3

Concentration: 1mg/ml

Form: Liquid

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Applications: WB (1:1000-2000)

Reactivity: Human (predicted: Sheep, Cow, Chicken, Dog, Horse)

Predicted MW: 86 kDa

Entrez Gene: 2261

Swiss Prot: P22607

Source: KLH conjugated synthetic peptide derived from human FGFR3: 501-600/801.


Purification: affinity purified by Protein A

Storage: 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Background: The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four cleavage sites are used; adrenocorticotrophin, essential for normal steroidogenesis and the maintenance of normal adrenal weight, and lipotropin beta are the major end products. In other tissues, including the hypothalamus, placenta, and epithelium, all cleavage sites may be used, giving rise to peptides with roles in pain and energy homeostasis, melanocyte stimulation, and immune modulation. These include several distinct melanotropins, lipotropins, and endorphins that are contained within the adrenocorticotrophin and beta-lipotropin peptides. Mutations in this gene have been associated with early onset obesity, adrenal insufficiency, and red hair pigmentation. Alternatively spliced transcript variants encoding the same protein have been described.

VALIDATION IMAGES



Sample: Lane 1: Human MCF-7 cell lysates Lane 2: Human HeLa cell lysates Primary: Anti-FGFR3 (bs-1301R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 86 kDa Observed band size: 100,115 kDa

PRODUCT SPECIFIC PUBLICATIONS

[IF=2.784] Ma C et al. Isolation and biological characteristic evaluation of a novel type of cartilage stem/progenitor cell derived from Small-tailed Han sheep embryos. *Int J Mol Med*. 2018 Jul;42(1):525-533. ICC ; Sheep . 29693133